

# Cingulo-Opercular Activity Predicts Subsequent Word Recognition

KI Vaden, SE Kuchinsky, SL Cute, JB Ahlstrom, JR Dubno, MA Eckert  
 Department of Otolaryngology-Head and Neck Surgery  
 Medical University of South Carolina, Charleston, SC



## Introduction

The engagement of attention systems is critical for successful communication, particularly when difficult listening conditions diminish speech intelligibility. Speech recognition requires more effort in noise and elicits increased activity in frontal cortices that are collectively referred to as the cingulo-opercular network [CO; 1,2].

The CO network responds to difficulty, response uncertainty, and errors across a range of experimental tasks [3]. Neuroimaging studies involving non-linguistic tasks have shown that CO activity also can predict behavioral adjustments on subsequent trials [4].

The goal of the current study was to assess the extent to which CO network activity predicts word recognition on subsequent trials in a normative sample of young adults.

### Predictions:

1. Elevated CO activity provides a word recognition benefit.
2. Coherent CO network engagement improves word recognition.

## Method

**Participants:** 18 young, normal hearing adults [20-38 years,  $m = 29.2$ ,  $sd = 5.8$ ; 10 females; native English speakers; right-handed distribution ( $m = 68.3$ ,  $sd = 50.3$ ; [7]). Mean pure tone thresholds (200 Hz to 8000 Hz) were less than 9.2 dB HL.


**Task:** Listen, then repeat the word aloud, or say "nope" if it was not recognizable.

**Stimuli:** 120 consonant-vowel-consonant words in multi-talker babble.


- Multi-talker babble: 82 dB SPL; speech: 85 dB SPL (60 words, +3 dB SNR).
- Multi-talker babble: 82 dB SPL; speech: 92 dB SPL (60 words, +10 dB SNR).
- Words were presented in alternating SNR-blocks of 4-6 trials, 60 trials per epoch.

**Word recognition:** Correct if the word was repeated exactly as it was presented.

### Trial Design



### Experiment Design



E-Prime presented scanner-synchronized stimuli with Sensimetrics piezoelectric insert earphones and recorded responses with a Resonance Technology microphone.

**fMRI:** 180 T2\*-weighted images (3 mm<sup>3</sup> voxels); TR = 8.6 sec; 25 min 48 sec.

**Structural MRI:** T1-weighted images (1 mm<sup>3</sup> voxels).

## Analysis

**Preprocessing.** Functional images were realigned, co-registered, and smoothed (8mm FWHM), detrended [5], and spatially normalized into the mean sample space derived from the T1-anatomical images [6].

General linear model and group-level SPM analyses were used to examine BOLD responses to babble, words and babble in each SNR condition, in addition to correct or incorrect word recognition trials.

A general linear mixed model (GLMM) analysis was performed to predict trial-level word recognition ( $W$ ) with normalized BOLD activity from the preceding trial in each voxel:  $W_t = SNR_t + BOLD_{t-1} + (1|SUB) + error$ .

Another GLMM analysis tested the association between the proportion of regions of interest (PROI) with elevated activity for each trial and next trial word recognition:  $W_t = SNR_t + PROI_{t-1} + (1|SUB) + error$ . Each participant's ROIs were defined independently of their BOLD time series by performing the voxel-level GLMM and leaving that participant out.


A voxel level threshold of  $Z = 3.09$ ,  $p = 0.001$  and cluster extent threshold = 20 voxels,  $p_{FWE} < 0.05$  were used for all fMRI results.

## Results


**Word Recognition:** poorer in the +3 dB SNR ( $66.1 \pm 7.6\%$ ) than in the +10 dB SNR ( $90.9 \pm 3.9\%$ ),  $Z = 11.34$ ,  $p < 0.001$ .

### BOLD Response

#### Word Recognition Errors




#### SNR Condition (+3 dB SNR > +10 dB SNR)



CO network activity increased following errors (red bar height denotes mean error for each trial) and during +3 dB SNR trials (dark blue bars) compared to +10 dB SNR trials (light blue bars).

### BOLD Associated with Improved Subsequent Word Recognition



#### Connectivity: Task

R.STS	APC	R.IFG	R.AIFO	L.IFG	DPC
0.38**	0.34**	0.43**	0.48**	0.51**	
	0.17**	0.64**	0.46**	0.46**	R.STS
		0.17**	0.35**	0.27**	APC
			0.51**	0.45**	R.IFG
				0.39**	R.AIFO

#### Connectivity: Rest

R.STS	APC	R.IFG	R.AIFO	L.IFG	DPC
0.30**	0.15	0.34**	0.40**	0.44**	
	0.12	0.56**	0.29**	0.30**	R.STS
		0.04	0.25**	0.12	APC
			0.31**	0.43**	R.IFG
				0.21**	R.AIFO

CO network activity increased before correct responses, after controlling for SNR effects on activity. Connectivity between CO regions increased during word recognition compared to rest epochs,  $Z = 4.93$ ,  $p < 0.001$ . Correct responses were more likely after more CO regions exhibited elevated activity,  $Z = 2.03$ ,  $p = 0.04$ .

## Conclusions

Elevated CO activity is frequently observed when speech recognition requires greater effort [1-2, 7-10].

Our results show that CO engagement increases the likelihood of correct word recognition on the next trial, similar to findings in visuospatial studies [4,11]. Furthermore, the trial-level connectivity analysis demonstrated that correct word recognition increased when all regions in the CO network exhibited elevated activity on the previous trial.

These findings are consistent with the premise that the CO network is important for adaptive control, including during word recognition.

### References

1. Eckert MA, Menon V, Walczak A, Ahlstrom J, Denelow S, Horwitz A, Dubno JR (2009) At the heart of the ventral attention system: the right anterior insula. *Hum Brain Mapp* 30:2530-2541.
2. Wild CJ, Yusuf A, Wilson DE, Peelle JE, Davis MH, Johnsrude IS (2012) Effortful listening: the processing of degraded speech depends critically on attention. *J Neurosci* 32:14010-14021.
3. Dosenbach NJF, Visscher KM, Palmer ED, Miezin FM, Wenger KK, Kang HC, Burgund ED, Grimes AL, Schlaggar BL, Petersen SE (2006) A core system for the implementation of task sets. *Neuron* 50:799-812.
4. Kerns JG, Cohen JD, MacDonald AW, Cho RY, Stenger VA, Carter CS (2004) Anterior cingulate conflict monitoring and adjustments in control. *Science* 303:1023-1026.
5. Macey PM, Macey KE, Kumar R, Harper RM (2004) A method for removal of global effects from fMRI time series. *Neuroimage* 22:360-366.
6. Avants BB, Tustison NJ, Song G (2011) Advanced normalization tools (www.slicer.org).
7. Sharp DJ, Scott SK, Mehta MA, Wise RJS (2006) The neural correlates of declining performance with age: evidence for age-related changes in cognitive control. *Cereb Cortex* 16:1739-1749.
8. Harris KC, Dubno JR, Keren NI, Ahlstrom JB, Eckert MA (2009) Speech recognition in younger and older adults: a dependency on low-level auditory cortex. *J Neurosci* 29:6078-6087.
9. Adank P, Davis MH, Hagoort P (2012) Neural dissociation in processing noise and accent in spoken language comprehension. *Neuropsychologia* 50:77-84.
10. Erb J, Henry MJ, Eisner F, Obleser J (2013) The brain dynamics of rapid perceptual adaptation to adverse listening conditions. *J Neurosci* 33:10688-10697.
11. Weissman DH, Roberts KC, Visscher KM, Woldorff MG (2006) The neural bases of momentary lapses in attention. *Nat Neuro* 9:971-978.

**Acknowledgements.** This work was supported by the National Institutes of Health / National Institute on Deafness and Other Communication Disorders (P50 DC 00422), MUSC Center for Advanced Imaging Research, South Carolina Clinical and Translational Research (SCTR) Institute, NIH/NICRR Grant number UL1 RR029882. This investigation was conducted in a facility constructed with support from Research Facilities Improvement Program (C06 RR14516) from the National Center for Research Resources, National Institutes of Health. We thank the study participants.